

Sexual Violence and Primary Health workers: Identification of training and collaborative needs

Fawole Olufunmilayo I.

Dept. of Epidemiology and Medical Statistics,
Faculty of Public Health, College of Medicine,
University of Ibadan, Nigeria.

SVRI Conference, J'burg, South Africa





July 2009



INTRODUCTION

- Sexual violence (SV) is a profound human rights violation. [1]
- SV impacts severely on the mental and physical health of survivors. [1]
- All women are vulnerable, irrespective of marital status, sexuality and age. [2]
- Rape is the most outrageous form of sexual violence. [1,3]
- Prevalence of rape in Nigeria ranges between 4 and 6%. [4&5]

- 
- Primary care personnel could be an ideal network to address SV, especially in low income countries where specialized manpower is few. [6]
 - In Nigeria, primary health care (PHC) facilities are strategically located, situated in both rural and urban areas and are well utilized by women. [7]
 - Most women see PHC providers as people with whom they can relate. [8]

- 
- Efforts to protect and help sexually abused women will remain futile without early identification and referral of such women by attending health care personnel. [9]
 - The knowledge, attitude and training of PHC workers and management practices of sexually violated women are presented.

METHODOLOGY

Study area: The study was conducted in the 33 local government areas (LGAS) of Oyo state.


Study sites : PHC facilities comprising of PHC, maternity centres, health post and dispensaries.

Study Design : Analytical cross-sectional survey.


Target population: Health workers who manage women at primary health care level.

Pilot study : Done on 20 PHC workers in LGA in Osun state.

Oyo State, Nigeria



• Total Popn : 4,587,764



■ 33 LGAs



- Ethical Clearance was obtained from the UI/UCH Institutional Review Board.
-

- Written informed consent was obtained from respondents.
- Data was collected with a 37 item semi-structured self administered questionnaire.
- 596 health workers from 104 health facilities in the 33 LGAs were interviewed.
- Data Management with SPSS 15.0 software.

RESULTS

Socio demographic characteristics

- Mean age of the respondents was 38.0 ± 8.16 years. Age range was 20-60 years.
- Years of work experience ranged from one to 37 years, with median of 12 years.


Table 1: Respondents Socio demographic Characteristics

Characteristic	Number Percent
Type of Health Facility	
PH centre	430(72.1)
Maternity centre	110(18.5)
Health post and dispensary	56(9.4)
Age (years)	
20-29	84(14.1)
30-39	244(40.9)
40-49	222(37.3)
>50	46(7.7)
Sex	
Male	64(21.5)
Female	234(78.5)
Cadre	
Community health officer	104(17.4)
Matron	100(16.7)
Nurses	122(20.4)
CH extension workers	222(37.2)
Others	48(8.1)
Years of experience	
1- 9	198(29.9)
10-19	210(35.2)
20-29	208 (34.9)

Knowledge

- 483(81.0%) gave correct examples of sexually violent acts.
- This was stated as rape by 524(87.9%).
- The major health consequences of SV stated were HIV(52%), depression (27%), STIs (12%), and unwanted pregnancy (9%).
- Mean knowledge score out of 8 statements was 4.1 ± 1.6 .

Figure 1: Major source of knowledge on SV



■ TV ■ Other Hws ■ Radio ■ School ■ Newspaper ■ Religious Ldr

Perceptions/Attitudes

	Statement	No	%
1	Women are partially to blame for act of SV	367	(61.6)
2	Children are partially to blame for act of SV	48	(8.0)
3	It is an intrusion into the patients private life to ask about SV	252	(42.3)
4	I am confident to ask women about SV	298	(50.0)
5	I believe I can help women who have been sexually abused	446	(74.8)
6	I am willing to enquire about SV from patients if trained	500	(83.9)

Perceptions/Attitudes

- 90% estimated that less than 0.1% of women in their practice experience SV.
- Mean attitude score out of 6 was 2.5 ± 1.9 .
- Although not significant, males were less likely to be knowledgeable (OR 0.67; 95% CI 0.96-2.94) and have good attitudes (OR 0.78; 95: CI 0.44-1.40) than females.


Table 2: Logistic Regression Analysis of Factors Affecting Hws Knowledge and Attitude

Profile	Knowledge		Attitude	
	OR	P value	AOR	P value
Age <40 yrs ≥40 yrs	1	0.72; 0.16	1	0.73; 0.32
Sex Female Male	1	0.67; 0.07	1	0.78; 0.41
Cadre Junior Senior	1	1.05; 0.83	1	1.90; 0.008 2.05; 0.007
Yrs work experience 1-9 10-19 20-29	1	0.71; 0.23 0.64; 0.13	1	0.93; 0.82 0.58; 0.11 0.34; 0.05

Screening

- Only 91 (16.2%) health workers had ever asked patients, if they were victims of SV.
- 13 (2.6%) had ever identified a sexually abused woman in the course of their practice.
- Lack of education and training was stated as a barrier to screening by almost 70%.


Figure 2: Barriers to Screening women for SV



Management Practices

- ~~o 65 (10.9%) had managed sexually violated women in the last 5 years.~~
- o Health workers main response included referral (60.4%), counseling (30.3%) and treatment (9.3%).

Figure 3: Reasons for the limited Quality of care of Survivors



Training

- Only 18.8% (112) had ever received any training on gender based violence.
- Health workers with previous training on violence were three times more likely to screen (AOR 2.66; 95%CI: 1.52-4.63).
- Younger aged health workers were more likely to have had training than their older colleagues (OR 0.80; 95%CI 0.47-1.38).

Table 3: Logistic Regression Analysis of Factors Affecting Screening and Training


Profile	Screening		Training	
	OR	P value	AOR	P value
Age <40 yrs ≥40 yrs	1	0.65; 0.08	1	0.80; 0.42
Sex Female Male	1	0.8; 0.52	1	1.08; 0.80
Cadre Junior Senior	1	1.87; 0.59	1	1.08; 0.77
Yrs of experience 1-9 10-19 20-29	1	0.62; 0.11 0.65; 0.15	1	0.45; 0.02 0.41; 0.02 0.59; 0.10 0.52; 0.12
Previous Training No Yes	1	2.75; 0.00	2.66; 0.001	

DISCUSSION


- ~~Hw did not sufficiently investigate the possibility of exposure to SV in their patients.~~
- Although, victims may not reveal history of abuse initially because of shame and stigma, after a few visits women often develop a trusting relationship in the PHC worker and feel more comfortable to talk about abuse. [9, 10]
- Hws grossly underestimated the magnitude of SV, suggesting misconception on the extent of the problem.


- This may affect effectiveness of care given as patients health problems will not be linked with prior or ongoing violence. [11]
-

- Hw were willing to assist survivors, however level of knowledge was inadequate and attitudes poor.
- Training of the health workers will be a first step at instituting change in identification and management of survivors.
- Hw need to be trained to be non judgmental or blaming, sympathetic and be confident. Receiving support, empathy, understanding and caring are critical in the management of SV. [12]

- 
- The training curriculum of health workers at primary care level needs to be reviewed.

 - Screening was not routinely done
 - Standard screening questionnaire should be developed to help PHC workers ask about abuse. [13]
 - Guidelines and policies need to be developed to guide SV screening, referral and social support for survivors.

- 
- Communication and collaborations between the PHC facilities and other tiers of the health system including other groups working on SV (police, judiciary, NGOs and FBOs) need to occur.
 - Survivors are at high risk of experiencing psychological conditions including PTSD, depression, and phobias, and are at increased risk for revictimization. [14, 15]
 - Receiving help early is critical, earlier disclosure of experiences of violence has been found to be helpful in lowering the rate of depression for survivors. [15]

- 
- **Strengths-** This study one of the first studies on the knowledge and screening practices of health workers on SV in Nigeria.
 - Secondly, hw at PHC level, an underserved group, were interviewed.
 - Finally, both rural and urban health facilities were involved in the survey.
 - **Limitations-** Results may not be generaliseable to secondary, tertiary and private health facilities in the country.
 - -Based on self report.
 - Larger national studies required.

CONCLUSION

- Most health workers were willing to discuss SV, but lacked fundamental knowledge and had poor attitudes.

- The need for urgent training of health workers is evident.
- In addition to training, school curriculum of health providers need to be reviewed.
- Cooperative networks should be developed with NGOs, health institutions and law enforcement authorities.
- Handbooks, guidelines and protocols to guide management of SV should be developed.
- In future, health providers in Nigeria and other low income countries should be expected to play more important roles in diagnosing SV and preventing future injury and death of women.



Acknowledgement

- We thank the Oyo state Local Government Civil Service Commission and State Ministry of Health for facilitating the study.
- The WHO Oyo state office for their support.
- Funds for the study was provided by the University, Senate Research grant 07000457

THANK YOU FOR LISTENING

